

Plate 1



(a) Seasonal headwater springs and dry stream course of Cowside Beck at c. SD 888692.



(b) Cowside Beck, from near the confluence of Lower Beck, looking towards Yew Cogar Scar.



(c) Cowside Beck gorge showing the stream flanked by shelving moss-covered limestone bedding. (O.L. Gilbert)



(d) Lower Cowside Beck, here following a meandering course above its confluence with the Skirfare.

Plate 2



(a) Darnbrook Beck; headwater stream at c. 580 m on blanket bog over drift-covered Millstone Grit on Fountains Fell.



(b) Darnbrook Beck; waterfall and tumbled mass of limestone blocks below the Middle Limestone at c. 500–520 m. (D. Hodgson)



(c) Darnbrook Beck; waterfall and ravine in massive limestone (lower Wensleydales), c. 450 m, SD 885714 (D. Hodgson)



(d) Darnbrook valley. Darnbrook House (white) bottom left, Darnbrook Fell and Fountains Fell (extreme left) on skyline.

Plate 3



(a) General view of Lower Beck, looking across Cowside Beck towards Dew Bottoms.



(c) Large tufa screen ('waterfall' on O.S. maps) on upper part of Waterfall Beck.



(b) Darnbrook; sump at head of intermittently dry stretch above farm. Note moss cover on boulders.



(d) Lower part of Waterfall Beck showing tufa deposition on waterfalls and in stream bed. (O.L. Gilbert)